

## Research Article



# *Aphloia theiformis* (Vahl.) Benn.: A Plant with Various Therapeutic Properties

Marie Carene Nancy Picot-Allain\*

Department of Health Sciences, University of Mauritius, Mauritius

**Received Date:** 11 October 2018

**Accepted Date:** 11 December 2018

**Published Date:** 20 December 2018

## Citation

Picot-Allain MCN (2018) *Aphloia theiformis* (Vahl.) Benn.: A Plant with Various Therapeutic Properties. J Soil Plant Biol 2018(1): 16-20.

Correspondence should be addressed to Marie Carene Nancy Picot-Allain, Mauritius  
E-mail: [marie.picot1@umail.uom.ac.mu](mailto:marie.picot1@umail.uom.ac.mu)

## Copyright

Copyright © 2018 Marie Carene Nancy Picot-Allain. This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and work is properly cited.

## Abstract

*Aphloia theiformis* (Vahl.) Benn. is an evergreen medicinal plant possessing multiple therapeutic virtues. Traditional use of the plant among folk populations of Madagascar, Comoros Island, Mascarenes Islands, Seychelles Archipelago, and tropical Africa was geared towards the management of dysentery, fever, rheumatism, ulcers, jaundice, gastrointestinal infections, skin infections, cataract, and diabetes mellitus. Multiple lines of evidence from the literature appraise the biological activities of *Aphloia theiformis*. Recent *in vitro* studies support the inhibitory action of *Aphloia theiformis* methanol leaves extract on key enzymes related to diabetes, obesity, hypertension, Alzheimer's disease, and kidney stone formation. Such findings have led to the identification of mangiferin. *Aphloia theiformis* was also reported to possess anti-malaria, anti-Chikungunya, and "anti-aging" properties. The present monograph tries to comprehensively establish the ethnopharmacological uses, biological activities, as well as the botanical description of *Aphloia theiformis*. It is expected that this comprehensive monograph will support further scientific studies in order to validate other traditional use of *Aphloia theiformis*, hence opening avenues for the development of novel therapeutic entities.

## Keywords

Indian Ocean Islands; Mangiferin; Medicinal Plant; Therapeutic Properties

## Introduction

A strong body of evidence generated by ethnobotanical studies praise the use of medicinal plants by cultures across the globe [1]. Indeed, herbal medicinal systems, practices, and knowledge have been transmitted from one generation to the next over the years [2]. It has been postulated that up to 80% of the population of developing countries still rely on traditional medicine for their primary health case, either because of cultural tradition or scarcity of modern treatment facilities [3]. A substantial number of scientific studies attest that there is currently a renewed interest in medicinal plants for the development of novel therapeutic agents geared towards the prevention and treatment of various pathologies.


*Aphloia theiformis* (Vahl.) Benn. is presently the only known flowering species of the Aphloiaceae family [4]. Ethnobotanical evidences substantiate the use of *A. theiformis* by folk cultures of Madagascar, Comoros Island, Mascarenes Islands, Seychelles Archipelago, and tropical Africa for the management of several human ailments [5]. The binomial nomenclature includes the epithet '*Aphloia*' which stems from the Latin word 'a' meaning 'without' and 'phloios' meaning 'bark/skin/flower', while '*theiformis*' comes from 'thea'

which means ‘tea’ and ‘formis’ stands for ‘form’ [5,6]. *A. theiformis* is commonly known as ‘Fandamane’ or ‘Bois goyave/gouyave’ in Mauritius, ‘Change écorce/Bois change écorce’ or ‘Gouyavier marron/Gouyave marron’ in Réunion Island, ‘Bois d’anémone’ in Rodrigues, ‘Bwa merl/Bois merle’ in Seychelles, ‘Voafotsy’, ‘Fandramanana’, or ‘Maramanana’ in Madagascar [7].

In an endeavor to valorise *A. theiformis*, this paper attempts to present recent findings on the biological properties of *A. theiformis* and aims at extending knowledge on its botanical description, ethnopharmacological uses, and phytochemical profile to the scientific community.

## Botanical Description

*A. theiformis* is an evergreen treelet reaching 10 to 15 m height (Figure 1A). The trunk, measuring 30 cm in diameter, is black to blackish brown, deciduous in patches; the underlying bark is smooth and pale brown. The young branches are reddish and striated. Mature leaves are variable in shape with narrow to broadly elliptic or narrowly oboval to oboval-elliptic blade. The leaf margin is denticulate to glandular-dentate and measure between 3-8 cm long and 1.5-4.5 cm wide (Figure 1B) [8]. Flowers are axillary, solitary or in bunch with white to pale yellow sepals measuring between 5-6 mm in diameter (Figure 1C). The fruits are sub-globular to ovoid-pyriform, turn white at maturity and measure 6-8 mm long and 4-6 mm in diameter (Figure 1D) [5].



**Figure 1:** A: *Aphloia theiformis* whole plant, B: *Aphloia theiformis* leaves, C: *Aphloia theiformis* flowers, D: Different plant parts of *Aphloia theiformis* [8] 1: flowering branch, 2: flower head, 3: developing fruit, 4&6: different types of young leaves, 5: mature leaf, 7: branch with male flower, 8: male flower, 9: fruit.

## Ethnopharmacological Uses

Literature search revealed the extensive use of *A. theiformis* in traditional medicine. *A. theiformis* has been used alone or in combination with other medicinal plants for the management of various diseases including dysentery, fever, rheumatism,

ulcers, jaundice, gastrointestinal infections, skin infections, cataract, decrease cholesterol level, diabetes, wound healing, as a diuretic, and anti-pyretic [5,9,10]. Table 1 summarises the preparation methods involving the use of *A. theiformis* for the management of different human ailments.

Scientific name of plants	Plant part	Indication	Method of preparation and/or administration	Source
<i>Aphloia theiformis</i> , <i>Terminalia bentzoe</i>	Leaves, bark	Dysentery	10-15 leaves of <i>Aphloia theiformis</i> , 5-6 pieces of bark of <i>Terminalia bentzoe</i> can be consumed 2-3 times daily	[5]
<i>Aphloia theiformis</i> , <i>Erythroxylum laurifolium</i> , <i>Cocos nucifera</i>	Leaves, bark, roots	Fever	Few leaves of <i>Aphloia theiformis</i> , bark of <i>Erythroxylum laurifolium</i> , roots of <i>Cocos nucifera</i> . Half cup can be consumed 3 times daily	[5]
<i>Aphloia theiformis</i> , <i>Antidesma madagascariense</i> , <i>Toddalia asiatica</i>	Leaves	Jaundice	Prepare decoction using leaves of <i>Aphloia theiformis</i> , <i>Antidesma madagascariense</i> , <i>Toddalia asiatica</i>	[6]
<i>Aphloia theiformis</i>	Leaves	Diuretic	An tea prepared using dried leaves of <i>Aphloia theiformis</i> (20 g for 1 L of water) can be consumed 3-4 times daily	[11]
<i>Aphloia theiformis</i> , <i>Combretum micranthum</i>	Leaves	Fever, jaundice, ulcer, rheumatism, gastrointestinal infections	–	[11]
<i>Aphloia theiformis</i>	Leaves	Stomach pains	Decoction prepared from leaves of <i>Aphloia theiformis</i>	[5]
<i>Aphloia theiformis</i>	Leaves, roots	Impetigo	Decoction prepared from leaves and roots of <i>Aphloia theiformis</i>	[5]
<i>Aphloia theiformis</i> , <i>Piper nigrum</i> , <i>Cinnamomum verum</i>	–	Refreshing	<i>Aphloia theiformis</i> mixed with <i>Piper nigrum</i> or <i>Cinnamomum verum</i> is consumed as refreshing drink	[5]
<i>Aphloia theiformis</i> , <i>Dracaena reflexa</i> , <i>Centella asiatica</i>	Leaves	Teething children	A decoction prepared from the leaves of <i>Aphloia theiformis</i> , <i>Dracaena reflexa</i> , and <i>Centella asiatica</i>	[5]
<i>Aphloia theiformis</i>	Roots	Skin infections	A root decoction of <i>Aphloia theiformis</i> is used to wash skin infections	[6]
<i>Aphloia theiformis</i>	Leaves	Nail infection	A leaf poultice is applied to infected nail	[5]
<i>Aphloia theiformis</i>	Roots	Laxative	Decoction is prepared from the roots of <i>Aphloia theiformis</i>	[5]
<i>Aphloia theiformis</i>	Bark	Chikungunya	Decoction is prepared from the bark of <i>Aphloia theiformis</i>	[12]
<i>Erythroxylum laurifolium</i> , <i>Aphloia theiformis</i> , <i>Cassia fistula</i> , <i>Antidesma madagascariense</i> , <i>Heimia myrtifolia</i>	Leaves, bark,	Lower back ache caused by kidney stones or urinary infections	Decoction prepared from <i>Erythroxylum laurifolium</i> stem bark, <i>Aphloia theiformis</i> leaves, <i>Cassia fistula</i> leaves, <i>Antidesma madagascariense</i> leaves, <i>Heimia myrtifolia</i> leaves. Three cups can be taken orally per day in the morning, at noon and at night.	[13]
<i>Aphloia theiformis</i>	Leaves		In traditional Malagasy medicine, an infusion of the leaves is used to treat fever and stomach aches	[14]


Table 1: Ethnopharmacological data on *Aphloia theiformis*.

## Major Chemical Constituents

Preliminary phytochemical screening of the crude methanol and aqueous extracts prepared from the leaves of *A. theiformis* indicated the presence of alkaloids, phenols, flavonoids, saponins, and anthraquinones. Tormentonic acid ester glucoside, 23-hydroxytormentonic acid ester glucoside, and 6 $\beta$ -hydroxytormentonic acid ester glucoside were isolated and identified from *A. theiformis* methanol leaves extract [15]. More recently, a group of authors reported the presence mangiferin (Figure 2), a xanthone glucoside, in *A. theiformis* methanol leaves extract [16].

The immunomodulatory and anti-inflammatory activities of *A. theiformis* collected in Comoros Island has been investigated. The authors reported that the phenolic rich fraction of *A. theiformis* exhibited immunosuppressive effect on lymphocyte, immunostimulant effect on monocytes and granulocytes at high doses, and anti-inflammatory-like effect *in vivo* [19]. The methanol bark extract reduced nitrite release from macrophages with an IC<sub>50</sub> value of 19  $\mu\text{g ml}^{-1}$  showing promising anti-inflammatory effects [20].

The solvent-free extract of *A. theiformis* aerial parts obtained from microwave extraction showed potent antiviral effect against ZIKV strains of Africa and Asia and clinical isolates



## Biological Activities

*In vitro* testing of the crude methanol extract revealed inhibitory action against enzymes related to diabetes (IC<sub>50</sub> 67.71, 55.20, 19.73  $\mu\text{g ml}^{-1}$  for  $\alpha$ -amylase,  $\alpha$ -glucosidase, and glycogen phosphorylase, respectively), obesity (IC<sub>50</sub> 939.97, 696.22  $\mu\text{g ml}^{-1}$  for pancreatic lipase and cholesterol esterase, respectively), Alzheimer's disease (IC<sub>50</sub> 186.04  $\mu\text{g ml}^{-1}$  for acetyl cholinesterase), hypertension (IC<sub>50</sub> 162.17  $\mu\text{g ml}^{-1}$  for angiotensin converting enzyme), and kidney stone formation (IC<sub>50</sub> 148.80  $\mu\text{g ml}^{-1}$  for urease). *In silico* molecular docking showed that mangiferin, identified in the crude methanol extract of *A. theiformis* leaves, docked to the afore-mentioned enzymes, showing  $\pi$ - $\pi$  interactions and hydrogen bonds mostly. The anti-diabetic properties of *A. theiformis*, documented from traditional use, can thus be attributed to the presence of mangiferin. Additionally, *A. theiformis* expressed antiglycation properties and retarded glucose diffusion *in vitro*. Besides, *A. theiformis* methanol leaves extract significantly ( $p < 0.05$ ) scavenged nitric oxide (IC<sub>50</sub> 37.05  $\mu\text{g ml}^{-1}$ ), 2, 2-diphenyl-2-picrylhydrazyl (IC<sub>50</sub> 64.42  $\mu\text{g ml}^{-1}$ ), and hypochlorous acid (IC<sub>50</sub> 41.24  $\mu\text{g ml}^{-1}$ ) radicals compared to ascorbic acid (IC<sub>50</sub> 77.65, 189.91 and 59.18  $\mu\text{g ml}^{-1}$  for respective assays) [18].

of dengue virus. The extract acted directly on viral particles, preventing attachment to cell surface and entry in host cells [4]. Recently, a group of researchers indicated that *A. theiformis* displayed high anti-Chikungunya activity, justifying its use during the Chikungunya outbreaks in Reunion Island in 2006 [11]. *A. theiformis* methanol bark extract (IC<sub>50</sub> 13.3  $\mu\text{g ml}^{-1}$ ) showed antiplasmodial activity against 3D7 chloroquine sensitive strain of *Plasmodium falciparum*, the most noxious *Plasmodium* species responsible for malaria in man [21]. The anti-microbial activity of *A. theiformis* methanol leaves extract against *Staphylococcus aureus*, *Salmonella enteritis*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Bacillus subtilis*, *Sclerotinia sclerotium*, and *Candida albicans* using the disc diffusion technique has also been reported [22]. The "anti-aging" and photo-protective properties of *A. theiformis* leaves of Madagascar have been attributed to the presence of xanthones, such as, mangiferin [10]. Mangiferin was reported to inhibit the activity of elastase (IC<sub>50</sub> 139.64  $\mu\text{M}$ ) and collagenase (IC<sub>50</sub> 253.57  $\mu\text{M}$ ), two enzymes targeted for the development of anti-aging agents [23]. As a consequence, pure extract of mangiferin (85-93% w/w) obtained from *A. theiformis* leaves of Madagascar has been developed for the cosmetic industry [14].

Cytotoxicity studies on human embryonic kidney cell line

HEK-293 using the iCELLigence system have showed that the crude methanol extract and active fractions ( $IC_{50}$  values >  $20 \mu\text{g ml}^{-1}$ ) of the leaves of *A. theiformis* exhibited negligible cytotoxic activity [17]. Besides, *A. theiformis* methanolic bark extract exhibited moderate cytotoxic activity against WI38 ( $IC_{50}$   $58.3 \mu\text{g ml}^{-1}$ ), WS1 ( $IC_{50}$   $61 \mu\text{g ml}^{-1}$ ) fibroblasts, and A-549 epithelial cells ( $IC_{50}$   $40 \mu\text{g ml}^{-1}$ ) [20,21].

## Conclusion

In an attempt to validate ethnopharmacological data, scientific evaluation is essential. Thus, this paper appraised the therapeutic potential of *A. theiformis* by providing an account of different studies undertaken to evaluate its biological activity. *A. theiformis* has proved to be a promising plant possessing potent biological activities such as anti-microbial, antioxidant, anti-diabetic, and immunomodulatory properties. Additionally, findings summarised here, highlight the therapeutic function of mangiferin, a phytochemical ubiquitously present in *A. theiformis* leaves, and substantiate the need for further focus on this lead molecule for the development of novel drugs. However, to further appraise the traditional use of *A. theiformis* further scientific *in vivo* evaluations are crucial.

## References

- Picot-Allain CMN (2018) Medicinal Plants: Novel Therapeutic Solutions for Diabetes. *Ann Diabetes Res* 2: 1009.
- Zengin G, Guler GO, Aktumsek A, Ceylan R, Picot CMN, et al. (2015) Enzyme Inhibitory Properties, Antioxidant Activities, and Phytochemical Profile of Three Medicinal Plants from Turkey. *Adv Pharmacol Sci* 2015: 01-08.
- New WHO guidelines to promote proper use of alternative medicines (2004) Media centre, News releases 2004, WHO, Switzerland.
- Clain E, Sinigaglia L, Koishi AC, Gorgette O, Gadea G, et al. (2018) Extract from *Aphloia theiformis*, an edible indigenous plant from Reunion Island, impairs Zika virus attachment to the host cell surface. *Sci Rep* 8.
- Gurib-Fakim A, Brendler T (2004) Medicinal and Aromatic Plants of Indian Ocean Islands, Madagascar, Comoros, Seychelles and Mascarenes. *Medpharm Sci Publication*, Germany. Pg no: 252-253.
- Gurib-Fakim A (2002) Mauritius through its Medicinal Plants. *Le Printemps Ltd*, Mauritius. Pg no: 108-110.
- Picot MCN, Mahomoodally MF (2017) Effects of *Aphloia theiformis* on key enzymes related to diabetes mellitus. *Pharm Biol* 55: 864-872.
- Antoine R, Brenan JPM, Mangenot G (1982) Plants-Mascarene Islands-Identification. In: *Flore des Mascareignes*, La Réunion, Maurice, Rodrigues. The Royal botanic gardens, Kew, UK. 1: 153-160.
- Mootoosamy A, Fawzi Mahomoodally M (2014) Ethnomedicinal application of native remedies used against diabetes and related complications in Mauritius. *J Ethnopharmacol* 151: 413-444.
- Danthu P, Lubrano C, Flavet L, Rahajanirina V, Behra O, et al. (2010) Biological factors influencing production of xanthones in *Aphloia theiformis*. *Chem Biodivers* 7: 140-150.
- Gurib-Fakim A (2008) *Toutes les plantes qui soignent*. Michel Lafon Editions, France. Pg no: 349-350.
- Ledoux A, Cao M, Jansen O, Lucia M, Campos PE, et al. (2018) Antiplasmodial, anti-chikungunya virus and antioxidant activities of 64 endemic plants from the Mascarene Islands. *Int J Antimicrob Agents* 52: 622-628.
- Sreekeesoon DP, Mahomoodally MF (2014) Ethnopharmacological analysis of medicinal plants and animals used in the treatment and management of pain in Mauritius. *J Ethnopharmacol* 157: 181-200.
- Giboulot J, Herimamy A, Solofonirina R, Maillet G, Loiseau A, et al. (2016) Development of an HPTLC-UV densitometry method for quantification of aphloïol in *Aphloia theiformis* leaf extract. *Planta Med* 82: 1-381.
- Gopalsamy N, Vargas D, Guého J, Ricaud C, Hostettmann K (1988) Saponins from leaves of *Aphloia theiformis*. *Phytochem* 27: 3593-3595.
- Picot MCN, Bender O, Atalay A, Zengin G, Loffredo L, et al. (2017) Multiple pharmacological targets, cytotoxicity, and phytochemical profile of *Aphloia theiformis* (Vahl.) Benn. *Biomed Pharmacother* 89: 342-350.
- Picot MCN, Zengin G, Mollica A, Stefanucci A, Carradori S, et al. (2017) In vitro and In silico Studies of Mangiferin from *Aphloia Theiformis* on Key Enzymes Linked to Diabetes Type 2 and Associated Complications. *Med Chem* 13: 633-640.
- Picot MCN (2018) Multiple pharmacological potential of selected medicinal plants relevant to diabetes type 2 and related complications. University of Mauritius, Mauritius.
- Hsoidrou S, Bousta D, Lachkar M, Said Hassane SO, El Youbi-Hamsas A, et al. (2014) Phytochemical screening, immunomodulatory and anti-inflammatory activities of the phenolic fraction of *Aphloia theiformis* (Vahl) Benn. plant used in Comorian traditional medicine. *Phytothérapie* 12: 354-359.
- Jonville M, Kodja H, Strasberg D, Pichette A, Ollivier E, et al. (2011) Antiplasmodial, anti-inflammatory and cytotoxic activities of various plant extracts from the Mascarene Archipelago. *J Ethnopharmacol* 136: 525-531.
- Jonville M, Kodja H, Humeau L, Fournel J, De Mol P, et al. (2008) Screening of medicinal plants from Reunion Island for antimalarial and cytotoxic activity. *J Ethnopharmacol* 120: 382-386.
- Rangasamy O, Raoelison G, Rakotoniriana FE, Cheuk K, Urverg-Ratsimamanga S, et al. (2007) Screening for anti-infective properties of several medicinal plants of the Mauritian flora. *J Ethnopharmacol* 109: 331-337.
- Ochocka R, Hering A, Stefanowicz-Hajduk J, Cal K, Barańska H (2017) The effect of mangiferin on skin: Penetration, permeation and inhibition of ECM enzymes. *PLoS One* 12: 0181542.